



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO9031



ADULT AMBULATORY INFUSION ORDER

Antibiotic Therapy

(Aminoglycosides, Daptomycin, & Glycopeptides)

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Page 1 of 4

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note**.
2. Monitor drug levels and adjust dose as necessary.
 - a. DAPTOmycin: draw Creatine Phosphokinase (CPK) - Plasma, Weekly. Monitor CPK more frequently in patients with recent prior or concomitant therapy with an HMG-CoA reductase inhibitor, unexplained CPK increases, or renal impairment
 - b. Vancomycin: draw trough level just before the 4th dose and once weekly.
 - c. Aminoglycosides: For daily dosing, draw random level 12 hours after the start of infusion and once weekly. For every 8-12 hour dosing, draw peak and trough weekly. Troughs are drawn just before the dose and peaks are drawn 30 minutes after the end of the dose.

NURSING ORDERS:

1. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

Aminoglycosides:

LABS:

- CBC with differential, every _____ (visit)(days)(weeks)(months) – Circle One
- CMP, every _____ (visit)(days)(weeks)(months) – Circle One
- Urine Dipstick w/o micro (10 dip), weekly during therapy**

Daily dosing

- Random _____ level, 12 hours post-dose, weekly during therapy**

Traditional dosing

- Peak _____ level, weekly during therapy**
- Trough _____ level, weekly during therapy**
- Labs already drawn. Date: _____

MEDICATION:

- amikacin _____ mg/kg = _____ mg in sodium chloride 0.9% 100 mL IV, over 30-60 minutes
- gentamicin _____ mg/kg = _____ mg in sodium chloride 0.9% 100 mL IV, over 30-60 minutes
- tobramycin _____ mg/kg = _____ mg in sodium chloride 0.9% 100 mL IV, over 20-60 minutes

Interval: (must check one)

- ONCE
- Daily x _____ doses
- Every _____ days x _____ doses



Oregon Health & Science University
Hospital and Clinics Provider's Orders

OHSU
Health

ADULT AMBULATORY INFUSION ORDER
Antibiotic Therapy
(Aminoglycosides, Daptomycin, & Glycopeptides)

Page 2 of 4

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

DAPTOmycin:

LABS:

- CBC with differential, every _____ (visit)(days)(weeks)(months) – Circle One
- CMP, every _____ (visit)(days)(weeks)(months) – Circle One
- CK, PLASMA, ONCE prior to therapy**
- CK, PLASMA, weekly during therapy**
- Labs already drawn. Date: _____

MEDICATION:

- DAPTOmycin _____ mg/kg = _____ mg in sodium chloride 0.9% 50 mL IV, over 30 minutes

Interval: (must check one)

- ONCE
- Daily x _____ doses
- Every _____ days x _____ doses

Dalbavancin:

LABS:

- CBC with differential, every _____ (visit)(days)(weeks)(months) – Circle One
- CMP, every _____ (visit)(days)(weeks)(months) – Circle One
- C-reactive protein, every _____ (visit)(days)(weeks)(months) – Circle One
- Labs already drawn. Date: _____

MEDICATION:

- Single dose regimen**

dalbavancin (DALVANCE) **1500 mg** in dextrose 5%, intravenous, ONCE, over 30 minutes
Interval: ONCE

- Two-dose regimen**

dalbavancin (DALVANCE) **1000 mg** in dextrose 5%, intravenous, ONCE, over 30 minutes
Interval: ONCE
&

dalbavancin (DALVANCE) **500 mg** in dextrose 5%, intravenous, ONCE, over 30 minutes
Interval: ONCE, 7 days after initial dose

- Other**

dalbavancin (DALVANCE) _____ mg in dextrose 5%, intravenous, ONCE, over 30 minutes
Interval: _____



Oregon Health & Science University
Hospital and Clinics Provider's Orders

OHSU
Health

ADULT AMBULATORY INFUSION ORDER
Antibiotic Therapy
(Aminoglycosides, Daptomycin, & Glycopeptides)

Page 3 of 4

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

Vancomycin:

LABS:

- CBC with differential, every _____ (visit)(days)(weeks)(months) – Circle One
- CMP, every _____ (visit)(days)(weeks)(months) – Circle One
- Vancomycin trough, weekly during therapy (first level prior to 4th dose)**
- Labs already drawn. Date: _____

MEDICATION:

- vancomycin 750 mg in sodium chloride 0.9% 150 mL IV
- vancomycin 1000 mg in sodium chloride 0.9% 250 mL IV
- vancomycin 1250 mg in sodium chloride 0.9% 250 mL IV
- vancomycin 1500 mg in sodium chloride 0.9% 300 mL IV
- vancomycin 25 mg/kg/day in sodium chloride 0.9% IV, ONCE over 24 hours, continuous infusion via CADD (OHSU only)
- vancomycin 30 mg/kg/day in sodium chloride 0.9% IV, ONCE over 24 hours, continuous infusion via CADD (OHSU only)

Infuse doses up to 1000 mg over at least 60 minutes and doses greater than 1000 mg over 120 minutes. Infusion rate not to exceed 17 mg/min

Interval: (must check one)

- ONCE
- Daily x _____ doses
- Every _____ days x _____ doses

FOR InfuSystem™ AMBULATORY PUMP USE (OHSU only; hook up at infusion location):

Duration:

- _____ days

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction



Oregon Health & Science University
Hospital and Clinics Provider's Orders

OHSU
Health

ADULT AMBULATORY INFUSION ORDER
Antibiotic Therapy
(Aminoglycosides, Daptomycin, & Glycopeptides)

Page 4 of 4

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Please check the appropriate box for the patient's preferred clinic location:

Hillsboro Medical Center
Infusion Services
364 SE 8th Ave, Medical Plaza Suite 108B
Hillsboro, OR 97123
Phone number: (503) 681-4124
Fax number: (503) 681-4120

Adventist Health Portland
Infusion Services
10123 SE Market St
Portland, OR 97216
Phone number: (503) 261-6631
Fax number: (503) 261-6756

Mid-Columbia Medical Center
Celilo Cancer Center
1800 E 19th St
The Dalles, OR 97058
Phone number: (541) 296-7585
Fax number: (541) 296-7610