

ADULT AMBULATORY INFUSION ORDER

Sodium Ferric Gluconate Complex
(FERRLECIT) Infusion

Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

# 

\*\*This plan will expire after 365 days at which time a new order will need to be placed\*\*

#### **GUIDELINES FOR ORDERING**

- 1. Send FACE SHEET and H&P or most recent chart note.
- 2. Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion. Labs drawn date:

## **NURSING ORDERS:**

- 1. TREATMENT PARAMETERS Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
- 2. Instruct patient to set follow up appointment with provider for follow up labs.
- 3. Monitor patient for signs and symptoms of hypotension during and following administration. Monitor patient for signs and symptoms of hypersensitivity during the infusion and for at least 30 minutes after completion of the infusion.
- 4. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

## **MEDICATIONS:**

sodium ferric gluconate complex (FERRLECIT) 125 mg in sodium chloride 0.9% 100 mL, intravenous, over 1 hour

## Interval:

Once		
Other:		

# AS NEEDED MEDICATIONS:

1. sodium chloride 0.9% bolus, intravenous, 500 mL, AS NEEDED x1 dose, for vein discomfort tolerability. Give concurrently with ferric gluconate



# Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

# Health Sodium Ferric Gluconate Complex (FERRLECIT) Infusion

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

## **HYPERSENSITIVITY MEDICATIONS:**

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the
  infusion and notify provider immediately. Administer emergency medications per the Treatment
  Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for
  symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following: I am responsible for the care of the patient (who is ide I hold an active, unrestricted license to practice medic that corresponds with state where you provide care to state if not Oregon);	cine in: ☐ Oregon ☐	(check box	
My physician license Number is #	f practice and authoriz		
Provider signature:	Date/Time:		
Printed Name:	Phone:	Fax:	
Please check the appropriate box for the patient's	preferred clinic loca	tion:	
☐ Hillsboro Medical Center Infusion Services 364 SE 8th Ave, Medical Plaza Suite 108B Hillsboro, OR 97123 Phone number: (503) 681-4124 Fax number: (503) 681-4120	□ Adventist Health Portland Infusion Services 10123 SE Market St Portland, OR 97216 Phone number: (503) 261-6631 Fax number: (503) 261-6756		
☐ Mid-Columbia Medical Center			

Phone number: (541) 296-7585 Fax number: (541) 296-7610

Celilo Cancer Center 1800 E 19th St

The Dalles, OR 97058