

# Oregon Health & Science University Hospital and Clinics Provider's Orders



ADULT AMBULATORY INFUSION ORDER
Ferric Derisomaltose (MONOFERRIC)
Infusion
Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

# ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

Weight:	kg	Height: _	cm
Allergies:			
Diagnosis Code:			
Treatment Start Date	):		Patient to follow up with provider on date:
**This plan will exp	oire afte	r 365 days	at which time a new order will need to be placed**
<b>GUIDELINES FOR</b>	ORDER	ING	
<ol> <li>Send FACE</li> </ol>	<b>SHEET</b>	and H&P	or most recent chart note.
<ol><li>Ferritin must</li></ol>	be obta	ined prior t	o start of treatment. Labs drawn date:

## **NURSING ORDERS:**

- TREATMENT PARAMETERS Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
- 2. Instruct patient to set follow up appointment with provider for follow up labs.
- 3. Monitor patient for signs and symptoms of hypersensitivity during the infusion and for at least 30 minutes after completion of the infusion.
- 4. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

#### **MEDICATIONS:**

# Ferric Derisomaltose (MONOFERRIC) dosing: (must check one)

- O For weight greater than or equal to 50 kg: 1,000 mg in sodium chloride 0.9% 100 mL, intravenous, ONCE, over 20 minutes
- O For weight less than 50 kg: 20 mg/kg in sodium chloride 0.9% 100 mL, intravenous, ONCE, over 20 minutes

### AS NEEDED MEDICATIONS:

1. sodium chloride 0.9% bolus, intravenous, 500 mL, AS NEEDED x1 dose, for vein discomfort. Give concurrently with ferric gluconate

#### HYPERSENSITIVITY MEDICATIONS:

- 1. NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction



# Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Ferric Derisomaltose
(MONOFERRIC) Infusion

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

		\ /		
By signing below, I represent the following: I am responsible for the care of the patient (who is I hold an active, unrestricted license to practice me that corresponds with state where you provide care state if not Oregon);	dicine in: ☐ Oregon	□ (ch	eck box Specify	
My physician license Number is # PRESCRIPTION); and I am acting within my scope medication described above for the patient identifies	e of practice and author	OMPLETED TO BE A VAL rized by law to order Infusion	<u>.ID</u> on of the	
Provider signature:	Date/Time:			
Printed Name:	Phone:	Fax:		
Please check the appropriate box for the patien  ☐ Hillsboro Medical Center	t's preferred clinic lo  ☐ Adventist Hea			
Infusion Services 364 SE 8th Ave, Medical Plaza Suite 108B		ket St		
Hillsboro, OR 97123  Phone number: (503) 681-4124  Fax number: (503) 681-4120	Portland, OR 9 Phone number Fax number: (	<del>: (503) 261-6631</del>		
☐ Mid-Columbia Medical Center				
Celilo Cancer Center 1800 E 19th St				
The Dalles, OR 97058  Phone number: (541) 296-7585				
Fax number: (541) 296-7565				