

*Generic substitute unless checked \square

	*Patient name: *DOB:
e	*Diagnosis and ICD-10 code:
	Allergies:
e	Outpatient admit: Series One time
	Code status: ☐ Full code ☐ DNR ☐ Medications only ☐ Other (specify)
	Vital signs: □ Per protocol □ Other (specify)
	Access port per protocol (using sterile technique, etc.)
	Frequency of port access: \square One time order \square Biweekly \square Weekly \square Monthly
	☐ Other (specify)
	Draw blood for labs: ☐ Yes ☐ No If yes, frequency of labs:
	Lab draws: □ CBC □ CMP □ Hgb and HCT □ PT □ Renal □ BMP
	☐ Other (specify)
	If not drawing blood for labs, check patency of line by drawing back to get blood return and then flushing with 20 mL. NS. Follow with 5 mL 100 units/mL Heparin prior to de-accessing port.
	If port flushes, but does not give blood return, try repositioning the patient (sitting up straighter, lying flat, ar above head, etc.)
	If port does not give blood or flush, use following protocol for Cathflo-Activase:
	 For patient > 30 kg: Instill 2 mg/mL into the occluded lumen
	 For patient < 30 kg: Instill volume equivalent to 100% the volume of the occluded lumen (up to 2 mg/mL)
	• May repeat the ordered dose after 120 minutes if blood still cannot be aspirated from the access device.
	Always flush port with 5 mL 100 units/mL Heparin prior to de-accessing.
	When finished, de-access per protocol.



Physician Order: Port Flush Adventist Health Tillamook 1000 Third St., Tillamook OR 97141 { Patient label }