

☑ Urine pregnancy test PRN			
Code Considerations for Patients NEW to SPRAVA ☐ F32.0 Major depressive disorder, single episode, m ☐ F32.1 Major depressive disorder, single episode, m ☐ F32.2 Major depressive disorder, single, severe w/o psychotic features ☐ F32.9 Major depressive disorder, single episode, u ☐ F33.0 Major depressive disorder, recurrent, mild ☐ F33.1 Major depressive disorder, recurrent, mode ☐ F33.2 Major depressive disorder, recurrent, severe w/o psychotic features ☐ R45.851 Suicidal ideations	nild Rece noderate	e Considerations for Patients Alreiving SPRAVATO®: 2.4 Major depressive disorder, single partial remission 2.5 Major depressive disorder, single full remission 3.41 Major depressive disorder, repartial remission 3.42 Major depressive disorder, refull remission	e episode, gle episode, current,
 □ Standard initiation for treatment-resistant dep Starting dose = 56mg intranasal Contact prescriber after first dose and advance - Date/Time dose advanced to 84mg: Give twice weekly for 4 weeks, then contact pr - Date/Time treatment beyond 4 weeks autho Then give once weekly for 4 weeks. Then give every other week. 	e dose to 84mg intra	Signature: treatment response.	
 □ Continue treatment for depression: Dose (check one) = □ 56mg / □ 84mg 			
 Previous treatment location and phone #: Continue treatment for major depressive disor Give (check one): ☐ 56mg / ☐ 84mg intranasa Give twice weekly to complete 4 weeks. How many doses of ketamine or esketamine had 	der with acute suic al.	idal ideation or behavior:	
 Contingency medication orders: Ondansetron 4mg ORAL x1 as needed for nausea. Promethazine 25mg ORAL or 12.5mg IM x1 as needed for he Acetaminophen 650mg ORAL x1 as needed for he Alprazolam 0.5mg ORAL x1 as needed for severe 	eeded for nausea, if eadache.	ondansetron not effective.	
*Physician signature *Denotes field that must be completed by healthcare wor	*Da	*Time	



Physician Order: SPRAVATO® Adventist Health Tillamook 1000 Third St., Tillamook OR 97141 { Patient label }