

Anifrolumab-Fnia (Saphnelo) Every 4 weeks

Anifrolumab-Fnia (Saphnelo) 300 mg every 4 weeks

Protocol Types

Infusion Treatment 1, Infusion Treatment 2, Infusion Treatment 3, Infusion Treatment 4

Patient name: _____ DOB: _____

Please include: ☐ Patient face sheet ☐ Insurance authorization (if other than Medicaid or non-Advantage Medicare)

Diagnosis: _____

Review: Every 358 days

Orders

	Interval	Defer Until	Duration
--	----------	-------------	----------

Labs

☒ hCG, serum, qualitative	Every 28 days		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
Inclusion conditions: Patient could become pregnant			
☒ CBC and differential	Every 28 days		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☒ Comprehensive metabolic panel	Every 28 days		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			

Nursing Orders

☒ Treatment conditions	Every 28 days		Until discontinued
Until discontinued Starting when released Until Specified			
Hold treatment and contact provider if patient has signs and symptoms of infection (fever, skin infection, need for antibiotics etc).			
☒ Nursing communication	Every 28 days		Until discontinued
Until discontinued Starting when released Until Specified			
Monitor for infusion-related reactions for 30 minutes after completion of the first infusion. If no previous infusion reactions, monitoring not required for subsequent doses. Monitoring recommended for previous infusion reactions, contact provider for guidance.			

Therapeutic Medications

☒ anifrolumab (Saphnelo) 300 mg in sodium chloride 0.9 % 100 mL IVPB	Every 28 days		Until discontinued
300 mg, intravenous, Administer over: 30 Minutes, Once, Starting when released			
Allow to reach room temp, then infuse using low-protein binding 0.2 or 0.22 micron in-line filter. Flush set with 25 mL NS after completion. Do not administer other meds through same line			

Emergency Medications

☒ Nursing communication	PRN		Until discontinued
STAT, Until discontinued Starting when released Until Specified			
If hypersensitivity reaction occurs, stop infusion IMMEDIATELY and CALL provider.			
☒ Vital Signs	PRN		Until discontinued

	Interval	Defer Until	Duration
STAT, Every 15 min Starting when released If a hypersensitivity reaction occurs, perform vital signs every 15 minutes for one hour, then every 30 minutes for 2 hours.			
☒ Nursing communication Once Starting when released If hypersensitivity reaction occurs: - Elevate head of bed 30 degrees and place patient's head at 30 degree angle if possible - Start 2 liters oxygen per nasal cannula and maintain airway - Change and maintain IV line with 1000 mL 0.9% Normal Saline	PRN		Until discontinued
☒ sodium chloride 0.9 % infusion 30 mL/hr, intravenous, Continuous PRN, KVO, Starting when released, for 1 day	PRN		Until discontinued
☒ albuterol 2.5 mg /3 mL (0.083 %) nebulizer solution 2.5 mg 2.5 mg, nebulization, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day	PRN		Until discontinued
☒ albuterol HFA 90 mcg/act inhaler 2 puff 2 puff, inhalation, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day SHAKE WELL	PRN		Until discontinued
☒ acetaminophen (Tylenol) tablet 650 mg 650 mg, oral, Once as needed, fever, GREATER THAN or EQUAL TO 38 C, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☒ diphenhydrAMINE (BENADryl) injection 25 mg 25 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☒ diphenhydrAMINE (BENADryl) injection 50 mg 50 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☒ hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus - OR - For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☒ EPINEPHrine (EpiPen;Adrenalin) injection solution 0.3 mg 0.3 mg, intramuscular, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued

Appointment Requests

☒ Infusion Appointment Request Status: Once, Expected: S, Sched. Duration: 60 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after	Every 28 days		Until discontinued
--	---------------	--	--------------------

*Provider signature

Provider printed name

*Date

Time

*Provider phone#

***Denotes field that must be completed by healthcare worker**