

Antibiotic Therapy - Dalbavancin Two-Dose Regimen

dalbavancin (DALVANCE) two dose regimen of 1000 mg once and 500 mg 7 days after initial dose

Protocol Types

Infusion Treatment 1, Infusion Treatment 2, Infusion Treatment 3, Infusion Treatment 4

Patient name: _____ DOB: _____

Please include: ☐ Patient face sheet ☐ Insurance authorization (if other than Medicaid or non-Advantage Medicare)

Diagnosis: _____

Review: Every 358 days

Orders

	Interval	Defer Until	Duration
Labs			
☐ CBC and differential Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine	Once		1 treatment
☐ Comprehensive metabolic panel Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine	Once		1 treatment
☐ C-reactive protein Status: Once, Expected: S, Lab Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine	Once		1 treatment
Therapeutic Medications			
☒ dextrose 5 % infusion 30 mL/hr, intravenous, Continuous, for 1 hour Flush line before and after dalbavancin administration with D5W ONLY.	Once		1 treatment
☒ dalbavancin (Dalvance) IV 1,000 mg 1,000 mg, intravenous, Once, Starting when released Flush line pre and post administration with D5W	Once		1 treatment
☒ dextrose 5 % infusion 30 mL/hr, intravenous, Continuous, for 1 hour Flush line before and after dalbavancin administration with D5W ONLY.	Once	S+7	1 treatment
☒ dalbavancin (Dalvance) IV 500 mg 500 mg, intravenous, Once, Starting when released Flush line pre and post administration with D5W	Once	S+7	1 treatment
Emergency Medications			
☒ Nursing communication STAT, Until discontinued Starting when released Until Specified If hypersensitivity reaction occurs, stop infusion IMMEDIATELY and CALL provider.	PRN		Until discontinued
☒ Vital Signs STAT, Every 15 min Starting when released If a hypersensitivity reaction occurs, perform vital signs every 15 minutes for one hour, then every 30 minutes for 2 hours.	PRN		Until discontinued
☒ Nursing communication	PRN		Until discontinued

	Interval	Defer Until	Duration
Once Starting when released			
If hypersensitivity reaction occurs:			
- Elevate head of bed 30 degrees and place patient's head at 30 degree angle if possible			
- Start 2 liters oxygen per nasal cannula and maintain airway			
- Change and maintain IV line with 1000 mL 0.9% Normal Saline			
☒ sodium chloride 0.9 % infusion	PRN		Until discontinued
30 mL/hr, intravenous, Continuous PRN, KVO, Starting when released, for 1 day			
☒ albuterol 2.5 mg /3 mL (0.083 %) nebulizer solution 2.5 mg	PRN		Until discontinued
2.5 mg, nebulization, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day			
☒ albuterol HFA 90 mcg/act inhaler 2 puff	PRN		Until discontinued
2 puff, inhalation, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day			
SHAKE WELL			
☒ acetaminophen (Tylenol) tablet 650 mg	PRN		Until discontinued
650 mg, oral, Once as needed, fever, GREATER THAN or EQUAL TO 38 C, Starting when released, for 1 dose			
For mild/moderate reaction: flushing, rash, pruritus			
☒ diphenhydramine (BENADryl) injection 25 mg	PRN		Until discontinued
25 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose			
For mild/moderate reaction: flushing, rash, pruritus			
☒ diphenhydramine (BENADryl) injection 50 mg	PRN		Until discontinued
50 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose			
For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			
☒ hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg	PRN		Until discontinued
100 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose			
For mild/moderate reaction: flushing, rash, pruritus			
- OR -			
For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			
☒ EPINEPHrine (EpiPen;Adrenalin) injection solution 0.3 mg	PRN		Until discontinued
0.3 mg, intramuscular, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose			
For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			

Appointment Requests

☒ Infusion Appointment Request	Once		1 treatment
Status: Once, Expected: S, Sched. Duration: 60 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after			
☒ Infusion Appointment Request	Once	S+7	1 treatment
Status: Once, Expected: S, Sched. Duration: 60 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after			

*Provider signature

Provider printed name

*Date

Time

*Provider phone#

*Denotes field that must be completed by healthcare worker