

Antibiotic Therapy - Fluoroquinolones, Macrolides, Lincosamides or Tetracyclines Daily

Fluoroquinolones (ciprofloxacin, levofloxacin), Macrolides (azithromycin), Lincosamides (clindamycin) or Tetracyclines (doxycycline) daily for 4 weeks

Protocol Types

Infusion Treatment 1, Infusion Treatment 2, Infusion Treatment 3, Infusion Treatment 4

Patient name: _____ DOB: _____

Please include: ☐ Patient face sheet ☐ Insurance authorization (if other than Medicaid or non-Advantage Medicare)

Diagnosis: _____

Review: Every 25 days

Orders

	Interval	Defer Until	Duration
Provider Ordering Guidelines			
☒ Provider communication	Every 1 day		Until discontinued
Until discontinued Starting when released Until Specified			
Intended duration of antibiotic therapy: *** days			

****ENSURE INTERVAL IS UPDATED TO REFLECT INTENDED DURATION****

Labs

☐ CBC and differential	1 time a week		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☐ Basic metabolic panel	1 time a week		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☐ C-reactive protein	1 time a week		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			

Therapeutic Medications

☐ AH THERAPEUTIC MEDICATIONS FLUOROQUINOLONES (CIPROFLOXACIN / LEVOFLOXACIN) (NO DEFAULT DOSE)			
☐ ciprofloxacin (Cipro) IV	Every 1 day		Until discontinued
intravenous, Once, Starting when released			
☐ levoFLOxacin (Levaquin) IV	Every 1 day		Until discontinued
intravenous, Once, Starting when released			
☐ AH THERAPEUTIC MEDICATIONS MACROLIDES (AZITHROMYCIN) (NO DEFAULT DOSE)			
☐ azithromycin (Zithromax) IV	Every 1 day		Until discontinued
intravenous, Administer over: 60 Minutes, Once, Starting when released			

	Interval	Defer Until	Duration
☐ AH THERAPEUTIC MEDICATIONS LINCOSAMIDES (CLINDAMYCIN) (NO DEFAULT DOSE)			
☐ clindamycin (Cleocin) IV intravenous, Once, Starting when released	Every 1 day		Until discontinued
☐ AH THERAPEUTIC MEDICATIONS TETRACYCLINES (DOXYCYCLINE) (NO DEFAULT DOSE)			
☐ doxycycline (Vibramycin) IVPB intravenous, Administer over: 1 Hours, Once, Starting when released	Every 1 day		Until discontinued
Emergency Medications			
☒ Nursing communication STAT, Until discontinued Starting when released Until Specified If hypersensitivity reaction occurs, stop infusion IMMEDIATELY and CALL provider.	PRN		Until discontinued
☒ Vital Signs STAT, Every 15 min Starting when released If a hypersensitivity reaction occurs, perform vital signs every 15 minutes for one hour, then every 30 minutes for 2 hours.	PRN		Until discontinued
☒ Nursing communication Once Starting when released If hypersensitivity reaction occurs: - Elevate head of bed 30 degrees and place patient's head at 30 degree angle if possible - Start 2 liters oxygen per nasal cannula and maintain airway - Change and maintain IV line with 1000 mL 0.9% Normal Saline	PRN		Until discontinued
☒ sodium chloride 0.9 % infusion 30 mL/hr, intravenous, Continuous PRN, KVO, Starting when released, for 1 day	PRN		Until discontinued
☒ albuterol 2.5 mg /3 mL (0.083 %) nebulizer solution 2.5 mg 2.5 mg, nebulization, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day	PRN		Until discontinued
☒ albuterol HFA 90 mcg/act inhaler 2 puff 2 puff, inhalation, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day SHAKE WELL	PRN		Until discontinued
☒ acetaminophen (Tylenol) tablet 650 mg 650 mg, oral, Once as needed, fever, GREATER THAN or EQUAL TO 38 C, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☒ diphenhydrAMINE (BENADryl) injection 25 mg 25 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☒ diphenhydrAMINE (BENADryl) injection 50 mg 50 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☒ hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus - OR - For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☒ EPINEPHrine (EpiPen;Adrenalin) injection solution 0.3 mg	PRN		Until discontinued

	Interval	Defer Until	Duration
0.3 mg, intramuscular, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			
Appointment Requests			
☒ Infusion Appointment Request	Every 1 day		Until discontinued
Status: Once, Expected: S, Sched. Duration: 120 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after			

*Provider signature	Provider printed name	*Date	Time
---------------------	-----------------------	-------	------

*Provider phone#

***Denotes field that must be completed by healthcare worker**