

ECT therapy plan for use in the ECT workflow.

Protocol Types

ECT

Patient name: _____ DOB: _____

Please include: ☐ Patient face sheet ☐ Insurance authorization (if other than Medicaid or non-Advantage Medicare)

Diagnosis: _____

Orders

	Interval	Defer Until	Duration
ECT Pre-procedure Orders			
☒ Insert peripheral IV Continuous Starting when released Until Specified	Every visit		Until discontinued
☒ Oxygen Therapy - Device: Nasal Cannula; Rate in liters per minute: 2 lpm Continuous First occurrence now Until Specified Keep SPO2% GREATER than or equal to: 92% Device: Nasal Cannula Rate in liters per minute: 2 lpm	Every visit		Until discontinued
ECT Pre-procedure Medications			
☒ glycopyrrolate (Robinul) injection 0.004 mcg/kg (Treatment Plan) 0.004 mcg/kg, intramuscular, Starting when released	Every visit		Until discontinued
ECT Pre-procedure Medications Anxiety			
☒ diphenhydrAMINE (BENADryl) capsule 25 mg 25 mg, oral, Once, Starting when released	Every visit		Until discontinued
☒ hydrOXYzine pamoate (Vistaril) capsule 25 mg 25 mg, oral, Once, Starting when released	Every visit		Until discontinued
ECT Pre-procedure Medications Nausea			
☒ ondansetron (Zofran) IV 4 mg 4 mg, intravenous, Once, Starting when released	Every visit		Until discontinued
ECT Treatment			
☒ Electroconvulsive therapy Once Starting when released	Every visit		Until discontinued
Appointment Requests			
☒ ECT Appointment Request Status: Future, Expected: S, Expires: S+366, Sched. Duration: 240 minutes, Sched. Tolerance: Schedule appointment at most 1 days before or at most 1 days after	Once		1 treatment
ECT Post-procedure Orders			
☒ Discontinue IV Continuous Starting when released Until Specified	Every visit		Until discontinued

	Interval	Defer Until	Duration
☒ Oxygen Therapy - Device: Nasal Cannula; Rate in liters per minute: 2 lpm Continuous First occurrence now Until Specified Keep SPO2% GREATER than or equal to: 92% Device: Nasal Cannula Rate in liters per minute: 2 lpm	Every visit		Until discontinued
ECT Post-procedure Medications Pain			
☒ acetaminophen (Tylenol) tablet 487.5 mg 500 mg, oral, Once, Starting when released	Every visit		Until discontinued
☒ ibuprofen (Motrin) tablet 400 mg 400 mg, oral, Once, Starting when released	Every visit		Until discontinued
☒ HYDROcodone-acetaminophen (Norco) 5-325 MG per tablet 1 tablet 1 tablet, oral, Once, Starting when released	Every visit		Until discontinued
ECT Post-procedure Medications Anxiety			
☒ QUetiapine (SEROquel) tablet 25 mg 25 mg, oral, Once, Starting when released If ordered by enteral tube route, hold tube feedings for 30 minutes before administration; flush with 25 mL of sterile water. After administration of this medication, follow with a 50 mL flush of sterile water. Restart tube feedings after drug administration as ordered.	Every visit		Until discontinued
☒ risperiDONE (RisperDAL) tablet 0.5 mg 0.5 mg, oral, Once, Starting when released	Every visit		Until discontinued
☒ haloperidol (Haldol) tablet 0.5 mg 0.5 mg, oral, Once, Starting when released	Every visit		Until discontinued
☒ diphenhydrAMINE (BENADryl) capsule 25 mg 25 mg, oral, Once, Starting when released	Every visit		Until discontinued
☒ hydroXYzine pamoate (Vistaril) capsule 25 mg 25 mg, oral, Once, Starting when released	Every visit		Until discontinued
ECT Post-procedure Medications Nausea			
☒ droperidol (Inapsine) 2.5 mg in dextrose 5 % 50 mL IVPB 2.5 mg, intravenous, Administer over: 15 Minutes, Once, Starting when released Prophylaxis dose = 0.015-0.075 mg/kg/dose, PONV is recommended 0.1 mg/kg.	Every visit		Until discontinued

*Provider signature

Provider printed name

*Date

Time

*Provider phone#

*Denotes field that must be completed by healthcare worker