

Protocol Types

Infusion Treatment 1, Infusion Treatment 2, Infusion Treatment 3, Infusion Treatment 4

Patient name: _____ DOB: _____

Please include: ☐ Patient face sheet ☐ Insurance authorization (if other than Medicaid or non-Advantage Medicare)

Diagnosis: _____

Review: Every 358 days

Orders

	Interval	Defer Until	Duration
Provider Communication			
☒ Provider communication	Every 84 days		Until discontinued
Until discontinued Starting when released Until Specified			
Monitor patients for new or worsening hypertension, Raynaud's phenomenon, and constipation with serious complications.			
Recommended dosage is 100 mg as an intravenous infusion every 3 months. Some patients may benefit from a dosage of 300 mg as an intravenous infusion every 3 months. Reassess efficacy after 6 months of therapy. Consider periodic evaluation of ongoing need for treatment and a trial off therapy after 12 to 18 months, as clinically appropriate.			
Therapeutic Medications			
☒ eptinezumab (Vyepti) 100 mg in sodium chloride 0.9 % 101 mL	Every 84 days		Until discontinued
100 mg, intravenous, Administer over: 30 Minutes, Once, Starting when released			
Use a 0.2 or 0.22 micron filter. Do not mix or administer with other meds. Following infusion, flush line with 20 mL NS			
Emergency Medications			
☒ Nursing communication	PRN		Until discontinued
STAT, Until discontinued Starting when released Until Specified			
If hypersensitivity reaction occurs, stop infusion IMMEDIATELY and CALL provider.			
☒ Vital Signs	PRN		Until discontinued
STAT, Every 15 min Starting when released			
If a hypersensitivity reaction occurs, perform vital signs every 15 minutes for one hour, then every 30 minutes for 2 hours.			
☒ Nursing communication	PRN		Until discontinued
Once Starting when released			
If hypersensitivity reaction occurs:			
- Elevate head of bed 30 degrees and place patient's head at 30 degree angle if possible			
- Start 2 liters oxygen per nasal cannula and maintain airway			
- Change and maintain IV line with 1000 mL 0.9% Normal Saline			
☒ sodium chloride 0.9 % infusion	PRN		Until discontinued
30 mL/hr, intravenous, Continuous PRN, KVO, Starting when released, for 1 day			
☒ albuterol 2.5 mg /3 mL (0.083 %) nebulizer solution 2.5 mg	PRN		Until discontinued

	Interval	Defer Until	Duration
2.5 mg, nebulization, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day			
☒ albuterol HFA 90 mcg/act inhaler 2 puff	PRN		Until discontinued
2 puff, inhalation, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day SHAKE WELL			
☒ acetaminophen (Tylenol) tablet 650 mg	PRN		Until discontinued
650 mg, oral, Once as needed, fever, GREATER THAN or EQUAL TO 38 C, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus			
☒ diphenhydrAMINE (BENADryl) injection 25 mg	PRN		Until discontinued
25 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus			
☒ diphenhydrAMINE (BENADryl) injection 50 mg	PRN		Until discontinued
50 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			
☒ hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg	PRN		Until discontinued
100 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus - OR - For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			
☒ EPINEPHrine (EpiPen;Adrenalin) injection solution 0.3 mg	PRN		Until discontinued
0.3 mg, intramuscular, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			

Appointment Requests

☒ Infusion Appointment Request	Every 84 days	Until discontinued
Status: Once, Expected: S, Sched. Duration: 90 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after		

*Provider signature	Provider printed name	*Date	Time
---------------------	-----------------------	-------	------

*Provider phone#

***Denotes field that must be completed by healthcare worker**