Adventist Health

Golimumab (Simponi Aria) Maintenance Every 8 Weeks

golimumab (Simponi Aria) 2 mg/kg every 8 weeks

Protocol Types

Infusion Treatment 1, Infusion Treatment 2, Infusion Treatment 3, Infusion Treatment 4

Patient name: _____ DOB: _____
Please include: □ Patient face sheet □ Insurance authorization (if other than Medicaid or non-Advantage Medicare)
Diagnosis: _____

Review: Every 358 days

Orders

	Interval	Defer Until	Duration
Provider Communication			
☑ Provider communication	Every 56 days		Until discontinued
Until discontinued Starting when released Until Specified Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. A Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). Please send results with order. If result is indeterminate, a follow up chest X-ray must be performed to rule out TB.			
Supplemental Labs			
☐ Hepatitis B Screening Labs			
☑ Hepatitis B Surface Antigen	Every 365 days		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☑ Hepatitis B Surface Antibody Quant	Every 365 days		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☑ Hepatitis B Core Antibody, Total	Every 365 days		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☐ TB Screening Labs			
☐ QUANTIFERON TB GOLD	Every 365 days		Until discontinued
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☐ TB Skin Test	Every 365 days		Until discontinued
Routine, Once Starting when released, Miscellaneous Release to patient: Immediate Point Of Care			
☐ Read PPD	Every 365 days		Until discontinued
Routine, Once Starting when released, Miscellaneous Release to patient: Immediate Point Of Care			
Labs			
☑ hCG, serum, qualitative	Every 56 days		Until discontinued

	Interval	Defer Until	Duration
Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine Inclusion conditions: Patient could become pregnant			
☑ CBC and differential Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine	Every 56 days		Until discontinued
☑ Comprehensive metabolic panel Status: Once, Expected: S, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine	Every 56 days		Until discontinued

Nursing Orders

☑ Treatment conditions Until discontinued Starting when released Until Specified Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.	Every 56 days		Until discontinued
☑ Treatment conditions Until discontinued Starting when released Until Specified Hold treatment and contact provider if patient has signs and symptoms of infection (fever, skin infection, need for antibiotics etc).	Every 56 days		Until discontinued
☑ Nursing communication Until discontinued Starting when released Until Specified Monitor and record vital signs, tolerance, and signs and symptoms of reactions prior to infusion and at the end of infusion.	Every 56 days		Until discontinued

Therapeutic Medications

☑ golimumab (Simponi Aria) injection 2 mg/kg 2 mg/kg, intravenous, Once, Starting when released	Every 56 days		Until discontinued

Medications

☑ acetaminophen (Tylenol) tablet 650 mg 650 mg, oral, Once as needed, headaches, fever, chills, or malaise, Starting when released, Until Discontinued	Every 56 days		Until discontinued
☑ diphenhydrAMINE (BENADryl) 25 mg 25 mg, oral, Once as needed, itching - 1st line, rash, Starting when released, Until Discontinued Give diphenhydramine if patient has transportation. Give either loratadine or diphenhydramine, not both.	Every 56 days		Until discontinued
☑ loratadine (Claritin) tablet 10 mg 10 mg, oral, Once as needed, For itching or rash if diphenhydramine is not given, Starting when released, Until Discontinued Give loratadine if patient does NOT have transportation. Give either loratadine or diphenhydramine, not both.	Every 56 days		Until discontinued

Emergency Medications

☑ Nursing communication STAT, Until discontinued Starting when released Until Specified If hypersensitivity reaction occurs, stop infusion IMMEDIATELY and CALL provider.	PRN		Until discontinued
☑ Vital Signs STAT, Every 15 min Starting when released If a hypersensitivity reaction occurs, perform vital signs every 15 minutes for one hour, then every 30 minutes for 2 hours.	PRN		Until discontinued
☑ Nursing communication Once Starting when released If hypersensitivity reaction occurs: - Elevate head of bed 30 degrees and place patient's head at 30 degree angle if possible - Start 2 liters oxygen per nasal cannula and maintain airway - Change and maintain IV line with 1000 mL 0.9% Normal Saline	PRN		Until discontinued
☑ sodium chloride 0.9 % infusion 30 mL/hr, intravenous, Continuous PRN, KVO, Starting when released, for 1 day	PRN		Until discontinued
☑ albuterol 2.5 mg /3 mL (0.083 %) nebulizer solution 2.5 mg	PRN		Until discontinued

	Interval	Defer Until	Duration
2.5 mg, nebulization, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day			
☑ albuterol HFA 90 mcg/act inhaler 2 puff 2 puff, inhalation, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day SHAKE WELL	PRN		Until discontinued
☑ acetaminophen (Tylenol) tablet 650 mg 650 mg, oral, Once as needed, fever, GREATER THAN or EQUAL TO 38 C, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☑ diphenhydrAMINE (BENADryl) injection 25 mg 25 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☑ diphenhydrAMINE (BENADryl) injection 50 mg 50 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☑ hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus - OR - For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☑ EPINEPHrine (EpiPen;Adrenalin) injection solution 0.3 mg 0.3 mg, intramuscular, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued

Appointment Requests

☑ Infusion Appointment Request Status: Once, Expected: S, Sched. Duration: 60 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after	Every 56 days		Until discontinued

*Provider signature Provider printed name *Date Time

*Provider phone#

*Denotes Field that must be completed by healthcare worker