

Protocol Types

Infusion Treatment 1, Infusion Treatment 2, Infusion Treatment 3, Infusion Treatment 4

Patient name: _____ DOB: _____

Please include: ☐ Patient face sheet ☐ Insurance authorization (if other than Medicaid or non-Advantage Medicare) Diagnosis: _____

Review: Every 161 days

Orders

	Interval	Defer Until	Duration
Provider Communication			
☒ Provider communication	Every 28 days		Until discontinued
Until discontinued Starting when released Until Specified			
Correct hypocalcemia before initiating treatment. Adequately supplement patients with calcium and vitamin D.			
Monitor serum calcium as clinically indicated.			

Labs

☒ CBC and differential	Every 84 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER)			
☒ Comprehensive metabolic panel	Every 84 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER)			
☒ PSA	Every 84 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER)			
☒ Testosterone (Adult Males)	Every 84 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER)			
☒ Vitamin D 25 hydroxy	Every 84 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER)			

Therapeutic Medications

☐ leuprolide (Lupron Depot) injection 7.5 mg	Every 28 days		Until discontinued
7.5 mg, intramuscular, Once, Starting when released			
Administer as a single injection into the gluteal area, anterior thigh, or deltoid. Give within 2 hours of preparation.			
HAZARDOUS - Handle with care			
☐ leuprolide (1-month) (Eligard) injection 7.5 mg	Every 28 days		Until discontinued
7.5 mg, subcutaneous, Once, Starting when released			
Give within 30 minutes of preparation. Administer to upper buttocks, upper or mid-abdomen. Avoid brawny or fibrous tissue, or areas with hair, nodules, lesions, or excessive pigmentation.			
HAZARDOUS - Handle with care			

Emergency Medications

☒ Nursing communication	PRN		Until discontinued
STAT, Until discontinued Starting when released Until Specified			
If hypersensitivity reaction occurs, stop infusion IMMEDIATELY and CALL provider.			

	Interval	Defer Until	Duration
☒ Vital Signs STAT, Every 15 min Starting when released If a hypersensitivity reaction occurs, perform vital signs every 15 minutes for one hour, then every 30 minutes for 2 hours.	PRN		Until discontinued
☒ Nursing communication Once Starting when released If hypersensitivity reaction occurs: - Elevate head of bed 30 degrees and place patient's head at 30 degree angle if possible - Start 2 liters oxygen per nasal cannula and maintain airway - Change and maintain IV line with 1000 mL 0.9% Normal Saline	PRN		Until discontinued
☒ sodium chloride 0.9 % infusion 30 mL/hr, intravenous, Continuous PRN, KVO, Starting when released, for 1 day	PRN		Until discontinued
☒ albuterol 2.5 mg /3 mL (0.083 %) nebulizer solution 2.5 mg 2.5 mg, nebulization, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day	PRN		Until discontinued
☒ albuterol HFA 90 mcg/act inhaler 2 puff 2 puff, inhalation, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day SHAKE WELL	PRN		Until discontinued
☒ acetaminophen (Tylenol) tablet 650 mg 650 mg, oral, Once as needed, fever, GREATER THAN or EQUAL TO 38 C, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☒ diphenhydrAMINE (BENADryl) injection 25 mg 25 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus	PRN		Until discontinued
☒ diphenhydrAMINE (BENADryl) injection 50 mg 50 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☒ hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus - OR - For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued
☒ EPINEPHrine (EpiPen;Adrenalin) injection solution 0.3 mg 0.3 mg, intramuscular, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)	PRN		Until discontinued

Appointment Requests

☒ Infusion Appointment Request Status: Future, Expected: S, Expires: S+365, Sched. Duration: 30 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after	Every 28 days		Until discontinued
--	---------------	--	--------------------

*Provider signature

Provider printed name

*Date

Time

*Provider phone#

***Denotes field that must be completed by healthcare worker**