

Vedolizumab 300 mg IV every 8 weeks

Protocol Types

Infusion Treatment 1, Infusion Treatment 2, Infusion Treatment 3, Infusion Treatment 4

Patient name: _____ DOB: _____

Please include: ☐ Patient face sheet ☐ Insurance authorization (if other than Medicaid or non-Advantage Medicare) Diagnosis: _____

Review: Every 180 days

Orders

	Interval	Defer Until	Duration
--	----------	-------------	----------

Provider Communication

☒ Provider communication	Every 56 days		Until discontinued
Until discontinued Starting when released Until Specified			
Treatment with vedolizumab should not be administered to patients with active TB infection.			

Patient should have regular monitoring for infection, malignancy, and liver abnormalities throughout therapy.

Supplemental Labs

☐ Hepatitis B Screening Labs			
☒ Hepatitis B Surface Antigen	Every 365 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☒ Hepatitis B Surface Antibody Quant	Every 365 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☒ Hepatitis B Core Antibody, Total	Every 365 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☐ TB Screening Labs			
☐ QUANTIFERON TB GOLD	Every 365 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☐ TB Skin Test	Every 365 days		Until discontinued
Routine, Once Starting when released, Miscellaneous			
Release to patient: Immediate			
Point Of Care			
☐ Read PPD	Every 365 days		Until discontinued
Routine, Once Starting when released, Miscellaneous			
Release to patient: Immediate			
Point Of Care			

Labs

☒ hCG, serum, qualitative	Every 56 days		Until discontinued
---	---------------	--	--------------------

	Interval	Defer Until	Duration
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
Inclusion conditions: Patient could become pregnant			
☒ CBC and differential	Every 56 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
☒ Comprehensive metabolic panel	Every 56 days		Until discontinued
Status: Future, Expected: S, Expires: S+365, Clinic Collect, Blood, Venous, Blood, AH INTERNAL LAB (BEAKER), Routine			
Therapeutic Medications			
☒ vedolizumab (Entyvio) 300 mg in sodium chloride 0.9 % 255 mL IVPB	Every 56 days		Until discontinued
300 mg, intravenous, Administer over: 30 Minutes, Once Administer as soon as possible after reconstitution			
Medications			
☒ acetaminophen (Tylenol) tablet 650 mg	PRN		Until discontinued
650 mg, oral, Once as needed, headaches, fever, chills, or malaise, Starting when released, Until Discontinued			
☒ diphenhydramine (BENADryl) 25 mg	PRN		Until discontinued
25 mg, oral, Once as needed, itching - 1st line, rash, Starting when released, Until Discontinued Give diphenhydramine if patient has transportation. Give either loratadine or diphenhydramine, not both.			
☒ loratadine (Claritin) tablet 10 mg	PRN		Until discontinued
10 mg, oral, Once as needed, For itching or rash if diphenhydramine is not given, Starting when released, Until Discontinued Give loratadine if patient does NOT have transportation. Give either loratadine or diphenhydramine, not both.			
Emergency Medications			
☒ Nursing communication	PRN		Until discontinued
STAT, Until discontinued Starting when released Until Specified If hypersensitivity reaction occurs, stop infusion IMMEDIATELY and CALL provider.			
☒ Vital Signs	PRN		Until discontinued
STAT, Every 15 min Starting when released If a hypersensitivity reaction occurs, perform vital signs every 15 minutes for one hour, then every 30 minutes for 2 hours.			
☒ Nursing communication	PRN		Until discontinued
Once Starting when released If hypersensitivity reaction occurs: - Elevate head of bed 30 degrees and place patient's head at 30 degree angle if possible - Start 2 liters oxygen per nasal cannula and maintain airway - Change and maintain IV line with 1000 mL 0.9% Normal Saline			
☒ sodium chloride 0.9 % infusion	PRN		Until discontinued
30 mL/hr, intravenous, Continuous PRN, KVO, Starting when released, for 1 day			
☒ albuterol 2.5 mg /3 mL (0.083 %) nebulizer solution 2.5 mg	PRN		Until discontinued
2.5 mg, nebulization, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day			
☒ albuterol HFA 90 mcg/act inhaler 2 puff	PRN		Until discontinued
2 puff, inhalation, Every 4 hours PRN, Hypersensitivity reaction, Starting when released, for 1 day SHAKE WELL			
☒ acetaminophen (Tylenol) tablet 650 mg	PRN		Until discontinued
650 mg, oral, Once as needed, fever, GREATER THAN or EQUAL TO 38 C, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus			
☒ diphenhydramine (BENADryl) injection 25 mg	PRN		Until discontinued

	Interval	Defer Until	Duration
25 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus			
☒ diphenhydrAMINE (BENADryl) injection 50 mg	PRN		Until discontinued
50 mg, intravenous, Administer over: 2 Minutes, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			
☒ hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg	PRN		Until discontinued
100 mg, intravenous, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For mild/moderate reaction: flushing, rash, pruritus - OR - For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			
☒ EPINEPHrine (EpiPen;Adrenalin) injection solution 0.3 mg	PRN		Until discontinued
0.3 mg, intramuscular, Once as needed, Hypersensitivity reaction, Starting when released, for 1 dose For severe reaction: respiratory distress, stridor, wheezing, hypotension (SBP LESS THAN 100 mm Hg)			

Appointment Requests

☒ Infusion Appointment Request	Every 56 days	Until discontinued
Status: Future, Expected: S, Expires: S+365, Sched. Duration: 60 minutes, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after		

*Provider signature	Provider printed name	*Date	Time
---------------------	-----------------------	-------	------

*Provider phone#

***Denotes field that must be completed by healthcare worker**