

ORDERS ARE IN EFFECT UNLESS CROSSED OUT.

Exceptions: Orders preceded by a box (\square) require a \checkmark to initiate order. Orders with blanks indicate additional information is needed.

	*Patient name:				*DOB:	
*Date	*Diagnosis: ☑ Osteoporosis M81.0					
	Allergies:					
*Time	Code status: ☐ Full code ☐ DNR ☐ Medications only ☐ Other (specify)					
	Draw serum creatinine and serum calcium up to six months prior to infusion for calculations, or on day of infusion with IV start.					
	Vascular Access: ☐ Port ☐ PICC ☐ CVC ☐ Start SL					
	If blood work meets criteria, give Reclast 5 mg in 100 mL NS IV over 30 minutes					
	 Serum calcium must be 8.4 mg/dL, if calcium less than 8.4, advise patient to take 600 mg calcium BID and recheck in one week. 					
	Creatinine clearance must be greater than 35 mL/min					
	Educate patient about recommendation that vitamin D intake should be 2000IU daily, and to keep hydrated.					
	Additional orders:					
	☐ Discharge patient when complete if stable					
	*Healthcare provider's signature:			*Date:	*Time:	
For use by Outpatient Therapy Services nurse below this line						
Ht	incm.	W: lb	kg.	Age	Gender: 🗆 Male 🗖 Female	SCr:
Calcium level:		If calcium less than 8.4, contact ordering physician, advise patient to take 600mg calcium BII and recheck in one week.				e 600mg calcium BID
Creatinine clearance:		Calculate using Cockroft-Gault formula — must be greater than 35 mL/min.				
Males			Females			
CrCl (m	L/min) = (140-age) >		$\frac{\text{in kg}}{\text{max}} = \frac{\text{CrCl (mL/min)} = (140-\text{age}) \times (\text{weight in kg})}{\text{max}} \times (0.85) = \frac{\text{crcl (mL/min)}}{\text{max}} \times (0.85) $			
	72 x serum crea	itinine (mg/dL)		72 x serum creatinine (mg/dL)		

*Denotes field that must be completed by healthcare worker

FAX to 503-815-7515

Physician Order Form: Reclast Infusion

* 2 3 7 *

Adventist Health Tillamook 1000 Third St., Tillamook OR 97141 { Patient label }